

RESEARCH ARTICLE

Open Access

Medicare and Medicaid enrollment and outside hospitalizations among HIV-infected and uninfected veterans engaged in VA care: a retrospective cohort study

Harry Chang^{1,2*}, Janet Tate², Amy C Justice^{2,3,4} and Michael E Ohl^{5,6,7}

Abstract

Background: Many veterans engaged in care with the Veterans Administration (VA) health system are also enrolled in Medicare and/or Medicaid and may receive care both inside and outside of the VA. Use of dual health systems has been associated with worse outcomes. Veterans with HIV may have different rates of Medicare and Medicaid enrollment and may be at greater risk of poor outcomes related to non-VA use. This study compares the frequency and factors associated with Medicare and/or Medicaid enrollment and non-VA use in an HIV-infected and uninfected population of veterans.

Methods: We used data from the VA and Center for Medicare & Medicaid Services from 2004 and 2005 to determine the frequency of Medicare and/or Medicaid enrollment among a cohort of HIV-infected and uninfected veterans engaged in VA care. We then restricted the cohort to veterans enrolled in fee-for-service (FFS) Medicare and/or Medicaid with at least one hospitalization and identified characteristics associated with non-VA hospital admissions.

Results: HIV-infected veterans had higher rates of Medicare and/or Medicaid enrollment than uninfected veterans (38% vs. 33%, $p < 0.01$), though the opposite was true when our sample was limited to veterans 65 years and older (53% vs. 70%, $p < 0.01$). Among veterans enrolled in the VA and FFS Medicare and/or Medicaid, veterans with HIV had greater illness severity and more frequent hospitalizations, but were less likely to be hospitalized outside the VA (48% vs. 54%, $p < 0.01$). HIV infection was associated with lower odds of outside hospitalization (OR = 0.76 [95% CI: 0.68, 0.85]).

Conclusions: Veterans with HIV have higher rates of Medicare and/or Medicaid enrollment, but lower odds of non-VA hospitalization. The VA integrated model of HIV care may discourage outside use among HIV-infected veterans.

Keywords: HIV, AIDS, Veterans, Medicare, Medicaid, Fragmentation

Background

A significant proportion of veterans enrolled with the Veterans Administration (VA) healthcare system are also enrolled in Medicare and/or Medicaid ("CMS enrollment"). An expansion of Medicaid coverage under the Patient Protection and Affordable Care Act of 2010 (ACA) will likely increase the proportion of CMS enrolled veterans [1]. Use

of non-VA health care ("CMS use") among veterans already engaged in VA care can lead to fragmented, inefficient, lower quality care and increased mortality [2-6].

Previous work has described CMS enrollment and identified factors that influence CMS use among VA enrolled veterans, particularly veterans 65 years and older that have age qualified for Medicare [7-9]. Less is known about CMS use among veterans under 65 years of age, who may qualify for Medicare through disability, or those enrolled in Medicaid. People living with HIV are more likely to qualify for Medicare through disability than age, and Medicare currently provides coverage for

* Correspondence: Harry.Chang@yale.edu

¹Yale School of Medicine, Yale University, 333 Cedar Street, New Haven, CT 06510, USA

²Veterans Aging Cohort Study, VA Connecticut Healthcare System, 950 Campbell Avenue, West Haven, CT 06516, USA

Full list of author information is available at the end of the article

approximately one-fifth of the people receiving HIV care in the U.S. [10]. Medicaid coverage for HIV varies on a state-to-state basis, but it currently covers about 50% of the people receiving HIV care in the U.S. [11]. Little is known about CMS enrollment among HIV-infected veterans, and the frequency and drivers of CMS use among veterans with both VA and CMS coverage.

HIV-infected veterans are an aging cohort vulnerable to adverse outcomes because of the complexity of their care needs, including multimorbidity, polypharmacy, mental health issues, and substance use [12-14]. The VA is the largest provider of care for persons with HIV in the United States, with over 20,000 veterans in care, and has invested significant resources to create an integrated HIV care model [15,16]. The integrated model for HIV care in VA emphasizes case management and access to coordinated medical and behavioral health care services, including mental health care, urgent care, substance abuse and on-site pharmacy services. Previous research has demonstrated that integrated HIV care at the VA can lead to higher rates of viral load suppression [14]. Given the high, and potentially expanding, rates of CMS enrollment in the HIV population, an understanding of Medicare and Medicaid enrollment and the frequency and drivers of CMS use among veterans with HIV is necessary to inform efforts to coordinate care.

The objectives of this paper were to: 1) determine the frequency of Medicare and/or Medicaid enrollment among HIV-infected veterans engaged in VA care, compared with uninfected veterans; 2) determine the frequency of non-VA hospital admissions among male, CMS enrolled HIV-infected and uninfected veterans with any hospitalizations; and 3) identify characteristics that predict non-VA hospital admission among male HIV-infected veterans enrolled in Medicare and/or Medicaid, compared with uninfected veterans. We hypothesized that CMS enrolled HIV-infected veterans would be less likely to use non-VA care than their uninfected counterparts.

Methods

This retrospective cohort study included two populations of veterans in VA care in the United States during 2004 and 2005: 1) HIV-infected veterans; and 2) age, race and facility-matched uninfected comparators.

We used data from the Veterans Aging Cohort Study (VACS), a prospective, observational cohort study of HIV infected veterans and age, race and site matched controls. VACS is made up of eight urban sites, but also includes a virtual cohort (VACS-VC), a nationally representative sample of HIV-infected veterans and matched uninfected controls identified through the VA electronic medical record (EMR) system and followed since 1996 [17,18]. Demographic data, medical diagnoses (based on


International Classification of Diseases, Ninth Revision, Clinical Modification [ICD-9-CM] codes), and laboratory results were extracted from VA EMR. Medicare and/or Medicaid enrollment and utilization for these veterans were extracted from data provided by the Center for Medicare & Medicaid Services (CMS). The study received approval from Human Investigations Committee at Yale University and the VA Connecticut Healthcare System and was granted a waiver of informed consent.

We started with a cohort that included all veterans enrolled in the VACS-VC prior to January 1, 2003 (cohort 1 in Figure 1) and engaged in VA care. For the purpose of this analysis we defined engaged in care as at least one inpatient or outpatient visit to a VA medical center in 2003. Using CMS enrollment data we determined the frequency of Medicare and/or Medicaid enrollment in this cohort. To study frequency and drivers of CMS use we limited our cohort of VA enrolled veterans to include only those with coverage for CMS use. We narrowed our sample to veterans that were CMS enrolled in the 2004 and 2005 calendar years and had at least one inpatient admission in either VA or non-VA hospitals during this same period. From this group we excluded those CMS enrolled for less than 18 months in the 24 month period, enrolled in HMO plans (as defined in the CMS data file), and enrolled in Medicaid in more than one state. Finally, we excluded female veterans, due to their low numbers in our final sample (cohort 2 in Figure 1). Our final sample consisted of 7,765 male HIV infected and uninfected veterans enrolled in FFS Medicare and/or Medicaid with at least one inpatient admission in 2004 and 2005. We used this cohort to determine the frequency and predictors of non-VA hospital admissions.

We created a series of variables to identify factors associated with non-VA hospital admissions among male veterans, enrolled in fee for service (FFS) Medicare and/or Medicaid that received inpatient care in 2004 and 2005 (cohort 2). Our dependent variable was a dichotomous indicator of any non-VA admissions, (vs. only VA inpatient admissions). Hospital admissions were identified by combining VA EMR data with CMS FFS reimbursement files.

We considered a number of independent variables we believed could influence use of non-VA inpatient care, including demographics, access to care, illness severity, and admission to a medical or surgical intensive care unit (ICU). Age and race data were taken from VA administrative databases. Age was constructed as a dichotomous variable, under age 65 and 65 years and older. Race was divided into three categorical variables: White, African American, and Other.

Patients' access to care was measured by three variables: enrollment in Medicare, Medicaid or both programs; distance to nearest VA tertiary care center treated as a



categorical variable: 0 to 9.9 miles, 10-29.9 miles, 30-59.9 miles, 60-119.9 miles, and greater than 120 miles [19]; and an indicator of whether veterans were subject to co-pays for VA care. Co-pay status was in turn determined using data on service connection, medical conditions, and an income-based means test for each veteran.

We used the CMS hierarchical condition categories (HCC) model to calculate risk scores [20] and used the scores as a measure of patients’ illness severity. The HCC model is a risk adjustment method employed by CMS to predict prospective payments for patients based on the previous year’s claims data, using patient demographics and major medical conditions, and has been applied to the VA population in the past for similar analyses [7]. Each year the average risk score is set at one, and we divided our HCC risk scores into a dichotomous variable: scores greater than one and scores less than or equal to one. We created a binary variable for HIV status, identified through VA administrative data. We also used VA administrative data to identify veterans that carried a diagnosis of a substance use problem (illicit substance abuse or dependence) using ICD-9CM codes and a previously validated algorithm available on the VACS website (vacohort.org).

We generated categorical variables for number of inpatient admissions (one vs. two or more) and the admission diagnosis (ICU vs. non-ICU admission) because we believe both could influence the likelihood of a veteran having a non-VA admission. Veterans with more frequent admissions have more “opportunities” to be hospitalized in a non-VA facility. Similarly, patients in critical condition, requiring admission to ICUs, may be more likely to be routed to a non-VA hospital. Though these variables do not reflect patient behavior, they may influence the likelihood of an admission outside the VA and, thus, we controlled for these factors in our model.

Analysis

We first examined veterans in the VACS-VC (cohort 1) to determine the frequency of CMS enrollment among veterans engaged in care. We stratified our analysis by age and HIV status, and used student t-tests to compare the frequency of CMS enrollment between HIV-infected and uninfected veterans. Next, we described the cohort of male, veterans enrolled in FFS Medicare and/or Medicaid with at least one inpatient admission (cohort 2) using the independent variables listed above. We stratified cohort

2 by HIV status and used chi-squared tests to compare the two sub-populations. Finally, we explored the association between these independent variables and non-VA hospital admissions using univariate and multivariable logistic regression. We created a multivariate model that included all patients in cohort 2, as well as separate models stratified by HIV status. All analyses were conducted using SAS version 9.2.

Results

Overall, HIV-infected veterans engaged in VA care were more likely than uninfected veterans to be enrolled in Medicare and/or Medicaid (38% vs. 33%, $p < 0.01$) (Table 1). While HIV-infected veterans under age 65 were more likely to be CMS enrolled than uninfected veterans, (36% vs. 29%, $p < 0.01$), HIV-infected veterans 65 years and older were less likely to be CMS enrolled (53% vs. 70%, $p < 0.01$). The majority of CMS enrolled veterans were enrolled in Medicare, but HIV-infected veterans were more likely than uninfected veterans to be enrolled in Medicaid, either alone or in combination with Medicare.

Among veterans enrolled in FFS Medicare and/or Medicaid with at least one inpatient admission (cohort 2) the frequency of HIV-infected veterans with CMS use was lower than the frequency among uninfected veterans (48% vs. 54%, $p < 0.01$) (Table 2). Compared to uninfected veterans, HIV-infected veterans were younger and lived closer to VA hospitals. A greater proportion of HIV-infected veterans was also of African American race, had an HCC score greater than 1 and carried a diagnosis of substance use problem.

HIV-infected veterans had lower odds of CMS use than uninfected veterans (OR = 0.78 [95% CI: 0.71, 0.85]) (Table 3). In addition, older veterans and veterans with higher HCC risk scores had greater odds of CMS use than their respective younger and healthier counterparts. Among HIV-infected veterans a substance abuse diagnosis was associated with increased non-VA use (OR = 1.34 [95% CI: 1.14, 1.58]). There was not a significant association between substance abuse and non-VA use in uninfected veterans (OR = 0.91 [95% CI: 0.80, 1.04]).

The association between HIV status and decreased odds of non-VA use remained in the multivariable model (OR = 0.76 [95% CI: 0.68, 0.85]); however, there was no longer an association between higher HCC risk score and increased non-VA use (Table 4). We posited that number of admissions, which had the greatest marginal effect on the multivariate model, was overwhelming the effects of other variables. When we removed number of admissions from the adjusted model (analysis not shown), both higher HCC risk score and African American race had significant associations with increased non-VA use (OR = 1.28 [95% CI: 1.13, 1.44] and OR = 1.18 [95% CI: 1.05, 1.33], respectively).

After stratifying by HIV statuses, the predictors of CMS use were similar in both the HIV-infected and uninfected models (Table 4).

Discussion

In our sample of veterans engaged in VA care, we found that HIV-infected veterans were more likely than their uninfected counterparts to be enrolled in Medicare and/

Table 1 Medicare and/or Medicaid enrollment among veterans enrolled in VACS-VC¹ in 2004

	All patients <i>n</i> (%) <i>n</i> = 91369	HIV+ <i>n</i> (%) <i>n</i> = 29909	HIV- <i>n</i> (%) <i>n</i> = 61460	<i>p</i> -value
Total Unique in VACS-VC				
Medicare Only	20964 (23)	6461 (22)	14503 (24)	<.0001
Medicaid Only	5220 (6)	2203 (7)	3017 (5)	<.0001
Medicare and Medicaid	5290 (6)	2618 (9)	2672 (4)	<.0001
Total Enrolled in CMS²	31474 (34)	11282 (38)	20192 (33)	<.0001
Under 65	<i>n</i> = 82835	<i>n</i> = 27193	<i>n</i> = 55642	
Medicare Only	16309 (20)	5334 (20)	10975 (20)	0.7111
Medicaid Only	5187 (6)	2190 (8)	2997 (5)	<.0001
Medicare and Medicaid	4475 (5)	2329 (9)	2146 (4)	<.0001
Total Enrolled in CMS	25971 (31)	9853 (36)	16118 (29)	<.0001
65 and over	<i>n</i> = 8534	<i>n</i> = 2716	<i>n</i> = 5818	
Medicare Only	4655 (55)	1127 (41)	3528 (61)	<.0001
Medicaid Only	33 (0)	13 (0)	20 (0)	0.3497
Medicare and Medicaid	815 (10)	289 (11)	526 (9)	0.0192
Total Enrolled in CMS	5503 (64)	1429 (53)	4074 (70)	<.0001

¹Veterans Aging Cohort Study – Virtual Cohort.

²Center for Medicare & Medicaid Service.

Table 2 CMS¹ enrolled veterans with 1 or more inpatient admissions in 2004 and 2005

	All patients n (%) n = 7765	HIV+ n (%) n = 2799	HIV- n (%) n = 4966	p-value
DEMOGRAPHICS				
Age				
<65	6319 (81)	2405 (86)	3914 (79)	<.0001
≥65	1446 (19)	394 (14)	1052 (21)	
Race				
White	3903 (50)	1343 (48)	2560 (52)	0.009
African American	3755 (48)	1413 (50)	2342 (47)	
Other	107 (2)	43 (2)	64 (1)	
Distance from VA				
0-9.9 miles	3253 (42)	1445 (52)	1808 (36)	<.0001
10-29.9 miles	1824 (24)	599 (21)	1225 (25)	
30-59.9 miles	1270 (16)	393 (14)	877 (18)	
60-119.9 miles	1192 (15)	310 (11)	882 (18)	
≥120 miles	224 (3)	52 (2)	172 (3)	
Co-pay status				
No Co-pay	7174 (95)	2592 (96)	4582 (95)	0.0997
Co-pay	352 (5)	112 (4)	240 (5)	
ENROLLMENT				
Medicare Only	6408 (83)	2088 (74)	4320 (87)	<.0001
Medicaid Only	469 (6)	274 (10)	195 (4)	
Medicare and Medicaid	888 (11)	437 (16)	451 (9)	
HEALTH				
HIV				
Positive	2799 (36)			
Negative	4966 (64)			
HCC Risk Score²				
<1	4129 (53)	1040 (37)	3089 (62)	<.0001
≥1	3636 (47)	1759 (63)	1877 (38)	
INPATIENT ADMISSIONS				
Number of Admissions				
1	3186 (41)	1091 (39)	2095 (42)	0.0058
≥2	4579 (59)	1708 (61)	2871 (58)	
ICU³ Admission				
Yes	1637 (21)	575 (21)	1062 (21)	0.3822
No	6128 (79)	2224 (79)	3904 (79)	
Substance Abuse				
Yes	1958 (25)	828 (30)	1130 (23)	<.0001
No	5807 (75)	1971 (70)	3836 (77)	
UTILIZATION				
VA ⁴ Admissions Only	3757 (48)	1467 (52)	2290 (46)	<.0001
Non-VA Admission	4008 (52)	1332 (48)	2676 (54)	

¹Center for Medicare & Medicaid Services.²Hierarchical Conditional Category Risk Score.³Intensive Care Unit.⁴Veterans Administration.

or Medicaid, but this difference varied by age. HIV-infected veterans under age 65 were more likely to be CMS enrolled, while HIV-infected veterans 65 years and older were much less likely to be CMS enrolled. Confirming our initial hypothesis, we also found that male, HIV-infected veterans enrolled in FFS Medicare and/or Medicaid were less likely to have an outside hospitalization. This was despite the fact that these veterans were generally sicker, more likely to have multiple admissions, and more likely to be enrolled in FFS Medicaid – independent risk factors for greater non-VA use.

We suspect that HIV-infected veterans were less likely to use non-VA hospitals because they feel more tied to the VA – possibly due to the integrated care and the different providers they see in VA infectious disease clinics or the desire for continuity with established caregivers for what is historically a stigmatized disease. The prescription drug coverage afforded veterans – and the high costs of antiretroviral therapy – may also play a significant role. While every state provides some coverage for antiretroviral drugs, the coverage varies substantially. Very few states cover as many of these drugs as are covered nationally within VA. This conclusion is also supported from patient self report data in VACS 8 in which 96% of those on ART reported getting all their antiretroviral medications through VA [17].

Our findings on frequency of CMS enrollment among veterans were not completely consistent with previous literature. Overall, we found lower rates of Medicare enrollment, but higher rates of enrollment in Medicaid [21]. This difference was particularly evident among veterans 65 years and older where Medicare enrollment has been reported to be greater than 90% [7,21]. This difference is likely driven primarily by our initial cohort (cohort 1), which included only patients who were already engaged in VA care, i.e. were seen at the VA at least once in the past year. We suspect that veterans already receiving care at the VA are less likely to seek alternative forms of coverage than all veterans ever enrolled in VA care. The difference may also be partially attributed to the greater share of HIV-infected veterans in our sample than in previously studied groups. VACS-VC is a nationally representative sample of HIV-infected veterans, not all veterans enrolled in VA care. As noted in the background, HIV-infected veterans differentially qualify for Medicare coverage, primarily through disability [10]. Similarly, we believe that comorbid and socioeconomic factors that contribute to higher rates of Medicaid enrollment among all persons living with HIV contributed to the higher rates of Medicaid enrollment in our sample of veterans. Among HIV infected and uninfected veterans, some of these changes may also be partially driven by a change in the population utilizing VA care since 1999 [8,21], when Shen, et al. reported

Table 3 Characteristics associated with non-VA inpatient admissions in 2004 and 2005, univariate analysis

	All patients <i>n</i> (%)	p-value	HIV+ <i>n</i> (%)	p-value	HIV- <i>n</i> (%)	p-value
DEMOGRAPHICS						
Age						
<65	Ref		Ref		Ref	
≥65	1.86 (1.65-2.09)	<.0001	1.39 (1.12-1.72)	0.0028	2.05 (1.77-2.36)	<.0001
Race						
White	Ref		Ref		Ref	
African American	0.96 (0.87-1.04)	0.3126	1.08 (0.93-1.26)	0.3031	0.90 (0.81-1.01)	0.0696
Other	3.40 (2.13-5.41)	<.0001	3.39 (1.70-6.79)	0.0006	3.58 (1.90-6.74)	<.0001
Distance from VA						
0-9.9 miles	Ref		Ref		Ref	
10-29.9 miles	1.30 (1.16-1.46)	<.0001	1.16 (0.96-1.40)	0.1297	1.37 (1.19-1.59)	<.0001
30-59.9 miles	1.75 (1.54-2.00)	<.0001	1.45 (1.16-1.81)	0.0012	1.91 (1.62-2.25)	<.0001
60-119.9 miles	2.55 (2.22-2.93)	<.0001	1.92 (1.49-2.46)	<.0001	2.82 (2.38-3.34)	<.0001
≥120 miles	2.41 (1.82-3.20)	<.0001	2.95 (1.62-5.37)	0.0004	2.26 (1.63-3.12)	<.0001
Co-pay status						
No Co-pay	Ref		Ref		Ref	
Co-pay	2.58 (2.04-3.27)	<.0001	1.92 (1.30-2.84)	0.001	3.01 (2.22-4.08)	<.0001
ENROLLMENT						
Medicare	Ref		Ref		Ref	
Medicaid	1.94 (1.60-2.36)	<.0001	2.51 (1.94-3.27)	<.0001	1.75 (1.30-2.36)	0.0003
Medicare and Medicaid	2.40 (2.07-2.79)	<.0001	2.95 (2.38-3.68)	<.0001	2.26 (1.83-2.80)	<.0001
HEALTH						
HIV						
Positive	0.78 (0.71-0.85)	<.0001				
Negative	Ref					
HCC Risk Score¹						
<1	Ref		Ref		Ref	
≥1	1.14 (1.04-1.25)	0.004	1.26 (1.08-1.46)	0.0039	1.21 (1.08-1.36)	0.0014
INPATIENT ADMISSIONS						
Number of Admissions						
1	Ref		Ref		Ref	
≥2	3.06 (2.79-3.37)	<.0001	4.29 (3.64-5.06)	<.0001	2.64 (2.35-2.97)	<.0001
ICU² Admission						
Yes	2.21 (1.97-2.47)	<.0001	2.00 (1.66-2.42)	<.0001	2.34 (2.02-2.70)	<.0001
No	Ref		Ref		Ref	
Substance Abuse						
Yes	1.04 (0.94-1.15)	0.4531	1.34 (1.14-1.58)	0.0004	0.91 (0.80-1.04)	0.1549
No	Ref		Ref		Ref	

¹Hierarchical Conditional Category Risk Score.²Intensive Care Unit.

their data, and our results from five years later, and may also reflect an evolving perception of VA care or increased Medicare cost sharing.

Our findings regarding predictors of non-VA hospitalization among CMS enrolled veterans engaged in VA

care were consistent with previous literature. In general, these studies have found older age, increased distance to the VA, greater cost-sharing and worse health to be associated with non-VA use [7,8,22,23]. Our results confirm these findings among HIV-infected veterans, and

Table 4 Characteristics associated with non-VA inpatient admissions in 2004 and 2005, multivariate analysis

	All patients <i>n</i> (%)	p-value	HIV+ <i>n</i> (%)	p-value	HIV- <i>n</i> (%)	p-value
DEMOGRAPHICS						
Age						
<65	Ref		Ref		Ref	
≥65	1.90 (1.65-2.17)	<.0001	1.61 (1.25-2.08)	0.0002	2.00 (1.70-2.35)	<.0001
Race						
White	Ref		Ref		Ref	
African American	1.05 (0.94-1.16)	0.4032	1.02 (0.86-1.22)	0.8293	1.07 (0.94-1.22)	0.3128
Other	3.17 (1.91-5.27)	<.0001	3.58 (1.63-7.85)	0.0015	2.98 (1.52-5.83)	0.0015
Distance from VA						
0-9.9 miles	Ref		Ref		Ref	
10-29.9 miles	1.51 (1.33-1.72)	<.0001	1.41 (1.13-1.75)	0.0022	1.59 (1.35-1.86)	<.0001
30-59.9 miles	2.08 (1.79-2.41)	<.0001	1.78 (1.38-2.30)	<.0001	2.25 (1.87-2.70)	<.0001
60-119.9 miles	3.42 (2.92-4.01)	<.0001	3.06 (2.29-4.09)	<.0001	3.62 (2.99-4.37)	<.0001
≥120 miles	3.06 (2.23-4.19)	<.0001	5.67 (2.90-11.06)	<.0001	2.61 (1.83-3.74)	<.0001
Co-pay status						
No Co-pay	Ref		Ref		Ref	
Co-pay	2.66 (2.05-3.46)	<.0001	2.61 (1.68-4.05)	<.0001	2.76 (1.99-3.84)	<.0001
ENROLLMENT						
Medicare	Ref		Ref		Ref	
Medicaid	2.83 (2.28-3.52)	<.0001	2.93 (2.17-3.94)	<.0001	2.57 (1.85-3.57)	<.0001
Medicare and Medicaid	2.50 (2.11-2.95)	<.0001	2.89 (2.27-3.69)	<.0001	2.17 (1.72-2.73)	<.0001
HEALTH						
HIV						
Positive	0.76 (0.68-0.85)	<.0001				
Negative	Ref					
HCC Risk Score¹						
<1	Ref		Ref		Ref	
≥1	0.99 (0.89-1.10)	0.8071	0.95 (0.79-1.14)	0.5937	1.01 (0.88-1.15)	0.9127
INPATIENT ADMISSIONS						
Number of Admissions						
1	Ref		Ref		Ref	
≥2	3.10 (2.79-3.45)	<.0001	4.45 (3.69-5.37)	<.0001	2.62 (2.30-2.98)	<.0001
ICU² Admission						
Yes	1.73 (1.52-1.96)	<.0001	1.42 (1.15-1.76)	0.0012	1.94 (1.65-2.27)	<.0001
No	Ref		Ref		Ref	
Substance Abuse						
Yes	1.16 (1.03-1.31)	0.0179	1.28 (1.05-1.55)	0.0138	1.08 (0.92-1.26)	0.3356
No	Ref		Ref		Ref	

¹Hierarchical Conditional Category Risk Score.²Intensive Care Unit.

provide guidance to VA care providers on patients that are more likely to receive non-VA care and thus benefit from care coordination efforts with outside providers.

Our results should be viewed in light of several limitations – most importantly, our limited access to data on

private insurance enrollment and utilization in the VA population. As a result of the lack of claims data for patients hospitalized under private insurance, some patients classified as having only VA admissions may actually have unaccounted for admissions to non-VA

hospitals. Moreover, CMS enrollment with private insurance may lead to significantly lower costs for veterans, influencing their choice of health care setting. The VACS does collect some survey data on private insurance enrollment. Among veterans in VACS8, 15% of HIV-infected veterans and 20% of uninfected veterans reported enrollment in private insurance (analysis not shown). Given the lower rates of supplemental insurance in the HIV-infected population, we believe our finding that HIV-infected veterans are less likely to have non-VA admissions likely still holds true. Another limitation is that inpatient admissions are more likely to be influenced by factors beyond patient preference. We tried to control for those variables in our analysis, but using inpatient admissions as our outcome may limit the conclusions that can be drawn from our results. We chose inpatient admissions because non-VA inpatient care places HIV-infected veterans at risk for adverse events related to poor coordination of care transitions between outside hospitals and VHA HIV clinics; however, choice in outpatient setting may be a better reflection of patient preferences and will be the subject of future studies. We also did not take into account pathways to CMS enrollment. Veterans that were enrolled in Medicare or Medicaid prior to engagement with the VA system may be more likely to have non-VA admissions; the opposite may be true of veterans that were seeking care with the VA prior to enrolling with Medicare and/or Medicaid. In addition, though most HIV infected people with Medicare qualify through disability, it is unknown if this is also true among veterans.

The heterogeneity of Medicare and/or Medicaid from state to state further limits our ability to generalize the data. Medicare Advantage plans, i.e. managed care plans provided by private insurers, are highly variable by design, and Medicaid benefits and eligibility vary from state to state. Moreover, to date only 26 of the 50 states have chosen to accept the ACA provision to expand Medicaid, making comparisons between states more difficult in the post-ACA era. Finally, our findings are limited by the age of our data. We restricted our analysis to 2004 and 2005, the latest years we had complete data available. However, we believe our analysis uses more recent data than similar studies analyzing CMS enrollment and CMS use.

Conclusion

Many veterans engaged in care in the VA are also enrolled in Medicare and/or Medicaid, including a significant proportion of veterans under age 65. Like the general population, the rates of CMS enrollment are higher among HIV infected veterans. However, among veterans enrolled in both VA and CMS care, HIV-infected veterans have lower odds of non-VA hospitalization despite having greater illness severity and more frequent hospitalizations than

uninfected veterans. The VA's integrated model of care used to treat HIV/AIDS may explain why HIV-infected veterans were more likely to seek care within the VA.

The impact of CMS use on veterans with HIV and other complex medical conditions requires further investigation. HIV-infected veterans have complicated medical histories and are likely to experience inefficient care and poor outcomes related to non-VA use. Research that provides evidence of the cost and health implications of CMS use may guide the VA in how aggressively to pursue and retain these patients in the VA system.

Future research should also explore why veterans with HIV were less likely than uninfected veterans to use non-VA care. If specific aspects of VA's integrated care programs for veterans with HIV, such as intensive case management, are found to contribute to lower likelihood of non-VA admission, then this would inform strategies to limit non-VA use among other veteran populations at high risk from fragmented care, such as veterans with other major chronic diseases or illnesses.

CMS enrollment is a valuable resource for veterans who face higher copayments or live long distances from VA medical centers; however, among some veterans, CMS enrollment may lead to inefficient care and poor health outcomes. As health coverage is further expanded efforts should be made to ensure that additional coverage is reaching the populations that can benefit the most. The VA and federal government should shape their policies to incentivize use of a single health system and develop procedures to ensure comprehensive exchange of information between health systems.

Abbreviations

HIV: Human immunodeficiency virus; AIDS: Acquired immunodeficiency disease syndrome; VA: Veterans Administration; CMS: Center for Medicare & Medicaid Services; ACA: Affordable Care Act; VACS-VC: Veterans Aging Cohort Study – Virtual Cohort; FFS: Fee for service; ICU: Intensive care unit; HCC: Hierarchical condition category; 95% CI: 95% Confidence interval; OR: Odds ratio.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

HC, JT, AJ, and MO conceived this study together and participated in its design. HC and JT conducted the data extraction. HC performed the statistical analyses with support from JT and MO. HC, JT, AJ, and MO drafted the manuscript. All authors read and approved the final manuscript.

Acknowledgements

The work reported here was supported by The Veterans Aging Cohort Study, with funding from National Institute on Alcohol Abuse and Alcoholism (U10 AA 13566) and VHA Public Health Strategic Health Care Group, and housed at the West Haven VAMC.

The views expressed in this article are those of the authors and do not necessarily represent the views of the Department of Veterans Affairs.

Author details

¹Yale School of Medicine, Yale University, 333 Cedar Street, New Haven, CT 06510, USA. ²Veterans Aging Cohort Study, VA Connecticut Healthcare System, 950 Campbell Avenue, West Haven, CT 06516, USA. ³Department of Internal

Medicine, Yale School of Medicine, 330 Cedar St, Boardman 110, P.O. Box 208056, New Haven, CT 06520, USA. ⁴Department of Public Health, Yale School of Public Health, 60 College Street, P.O. Box 208034, New Haven, CT 06520, USA. ⁵VA Office of Rural Health (ORH), Midwest Rural Health Resource Center, Iowa City VAMC, 601 Highway 6 West (152), Iowa City, IA 52246, USA. ⁶Center for Comprehensive Access and Delivery Research and Evaluation (CADRE) at the Iowa City VA Medical Center, Mailstop 152, Iowa City VAMC, Iowa City, IA 52246, USA. ⁷Department of Internal Medicine, University of Iowa Carver College of Medicine, 200 Hawkins Drive, Iowa City, IA 52242, USA.

Received: 4 June 2014 Accepted: 6 January 2015

Published online: 22 January 2015

References

- Haley JK, Kenney GM. Uninsured veterans and family members: State and National Estimates of expanded medicaid eligibility under the ACA. In: Timely analysis of immediate health policy issues. Urban Institute; 2013.
- Trivedi AN, Grebla RC, Jiang L, Yoon J, Mor V, Kizer KW. Duplicate federal payments for dual enrollees in Medicare advantage plans and the Veterans affairs health care system. *JAMA*. 2012;308(1):67–72.
- Tarlov E, Lee TA, Weichle TW, Durazo-Arvizu R, Zhang Q, Perrin R, et al. Reduced overall and event-free survival among colon cancer patients using dual system care. *Cancer Epidemiol Biomarkers Prev*. 2012;21(12):2231–41.
- Wolinsky FD, An H, Liu L, Miller TR, Rosenthal GE. Exploring the association of dual use of the VHA and Medicare with mortality: separating the contributions of inpatient and outpatient services. *BMC Health Serv Res*. 2007;7:70.
- Wolinsky FD, Miller TR, An H, Brezinski PR, Vaughn TE, Rosenthal GE. Dual use of Medicare and the Veterans Health Administration: are there adverse health outcomes? *BMC Health Serv Res*. 2006;6:131.
- Helmer D, Sambamoorthi U, Shen Y, Tseng CL, Rajan M, Tiwari A, et al. Opting out of an integrated healthcare system: dual-system use is associated with poorer glycemic control in veterans with diabetes. *Prim Care Diabetes*. 2008;2(2):73–80.
- Hynes DM, Koelling K, Stroupe K, Arnold N, Mallin K, Sohn MW, et al. Veterans' access to and use of Medicare and Veterans Affairs health care. *Med Care*. 2007;45(3):214–23.
- Humensky J, Carretta H, De Groot K, Brown MM, Tarlov E, Hynes DM. Service utilization of veterans dually eligible for VA and medicare fee-for-service: 1999–2004. *Medicare Medicaid Res Rev*. 2012;2(3). doi:10.5600/mmrr.002.03.a06.
- Liu CF, Manning WG, Burgess JF, Sharp N, Perkins M, Maciejewski ML. Reliance on Veterans Affairs outpatient care by Medicare-eligible veterans. *Med Care*. 2011;49(10):911–7.
- Medicare and HIV/AIDS: fact sheet. In: the Henry J. Kaiser family foundation 2009. Last accessed 6 October 2014. http://kaiserfamilyfoundation.files.wordpress.com/2013/01/7171_04.pdf.
- Medicaid and HIV/AIDS: a national analysis. In: The Henry J. Kaiser family foundation October 2011. Last accessed 6 October 2014. <http://kff.org/hivaids/report/medicaid-and-hiv-a-national-analysis/>.
- Womack JA, Goulet JL, Gibert C, Brandt CA, Skanderson M, Gulanski B, et al. Physiologic frailty and fragility fracture in HIV-infected male veterans. *Clin Infect Dis*. 2013;56(10):1498–504.
- Justice AC, Braithwaite RS. Lessons learned from the first wave of aging with HIV. *AIDS*. 2012;26 Suppl 1:S11–8.
- Hoang T, Goetz MB, Yano EM, Rossman B, Anaya HD, Knapp H, et al. The impact of integrated HIV care on patient health outcomes. *Med Care*. 2009;47(5):560–7.
- Yano EM, Asch SM, Phillips B, Anaya H, Bowman C, Chang S, et al. Organization and management of care for military veterans with human immunodeficiency virus/acquired immunodeficiency syndrome in Department of Veterans Affairs Medical Centers. *Mil Med*. 2005;170(11):952–9.
- The State of Care for Veterans with HIV/AIDS: 2011 Summary Report, D.o.V. Affairs, Editor 2012. Last accessed 6 October 2014. <http://www.hiv.va.gov/pdf/VA2011-HIVSummaryRpt.pdf>.
- Justice AC, Dombrowski E, Conigliaro J, Fultz SL, Gibson D, Madenwald T, et al. Veterans Aging Cohort Study (VACS): overview and description. *Med Care*. 2006;44(8 Suppl 2):S13–24.
- Fultz SL, Skanderson M, Mole LA, Gandhi N, Bryant K, Crystal S, et al. Development and verification of a "virtual" cohort using the National VA Health Information System. *Med Care*. 2006;44(8 Suppl 2):S25–30.
- Bliss RL, Katz JN, Wright EA, Losina E. Estimating proximity to care: are straight line and zipcode centroid distances acceptable proxy measures? *Med Care*. 2012;50(1):99–106.
- Evans MA. Evaluation of the CMS-HCC Risk Adjustment Model, C.f.M.M. Services, Editor. 2011.
- Shen Y, Hendricks A, Wang F, Kazis LE. VHA enrollees' health care coverage and use of care. *Med Care Res Rev*. 2003;60(2):253–67.
- Petersen LA, Byrne MM, Daw CN, Hasche J, Reis B, Pietz K. Relationship between clinical conditions and use of Veterans Affairs health care among Medicare-enrolled veterans. *Health Serv Res*. 2010;45(3):762–91.
- Shen Y, Findley PA, Maney M, Pogach L, Crystal S, Rajan M, et al. Department of Veterans Affairs-Medicare dual beneficiaries with stroke: where do they get care? *J Rehabil Res Dev*. 2008;45(1):43–5.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

