

RESEARCH

Open Access



Design and evaluation of an educational mobile program for liver transplant patients

Mostafa Langarizadeh¹, Fateme Moghbeli², Shamim Ahmadi¹, Mohammad Hossein Langarizadeh³, Mohammadjavad Sayadi⁴, Fatemeh Sarpourian⁵ and Seyed Ali Fatemi Aghda^{1,6*}

Abstract

Background Liver transplantation, the last treatment for advanced liver failure, necessitates patient education due to its wide range of complications and subsequent disabilities. The present study was development-applied research and aimed to design a mobile-based educational program to provide liver transplant patients with critical health information.

Methods In the first phase of the study, the crucial educational components were collected from the literature and organized in the form of a questionnaire using library studies and available global guidelines. The validity and reliability of this researcher-made questionnaire were confirmed by a panel of experts ($n = 15$), including gastroenterologists and liver specialists working in the Motahari liver clinic and AbuAli Sina Hospital in Shiraz. The application was designed followed by analyzing the data gathered from the first phase. To evaluate the mobile phone program's usability, to evaluate the application, 30 liver transplant patients were randomly selected.

Results Most educational components covered in the questionnaire were deemed necessary by experts in the first phase. As a result, the educational contents were classified under 10 categories. The application had a good level of usability since the participants' satisfaction score was 8.1 (out of 9 points).

Conclusions Due to the increase in liver transplantation and the use of mobile phones, applications increase the patient's role in their health, and their awareness. It also leads to a better interaction and follow-up of the patient, the treatment staff of the medical centers.

Keywords Liver transplant, Educational needs, Self-care, Usability evaluation, Mobile phone health (mHealth)

*Correspondence:

Seyed Ali Fatemi Aghda
afatamy@yahoo.com

¹ Department of Health Information Management, School of Health Management and Information Sciences, Iran University of Medical Sciences, Tehran, Iran

² Department of Health Information Technology, Varastegan Institute for Medical Sciences, Mashhad, Iran

³ Department of Computer Sciences, Faculty of Mathematics Sciences, Shahid Beheshti University, Tehran, Iran

⁴ Department of Computer Engineering, Technical and Vocational University (TVU), Tehran, Iran

⁵ Department of Health Information Technology, School of Management and Medical Informatics, Shiraz University of Medical Sciences, Shiraz, Iran

⁶ Research Center for Health Technology Assessment and Medical Informatics, School of Public Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Introduction

Nowadays, the prevalence of chronic diseases is increasing at an exponential rate. This not only imposes negative impacts on people's quality of life but also increases health care costs. Chronic liver disease, as a serious progressive condition of liver function [1], is one of the main global causes of death causing a wide range of diseases from different or unknown sources. This disease has been directly responsible for two to four% of all annual deaths throughout the world [2]. The literature is replete with studies conducted on the causes of advanced chronic liver failure in different parts of the world, according to the patients' geographic region, race, lifestyle, age, and other variables. The results showed a discrepancy in findings regarding common causes of the disease in developing and high-income countries [3–7].

Liver transplantation is known as the last treatment for patients with chronic liver failure. In the last decade, the number of people waiting to receive a liver transplant has been increasing [8]. Although patients who undergo liver transplantation seek a better quality of life in the long term, they need intensive medical care after the operation that exposes them to various challenges, such as physical problems, fatigue, activity limitation, poor sleep, mental problems, as well as increased levels of stress, anxiety, and depression [9–11].

Lack of adherence to medications in transplanted patients, reflexes a serious problem leading to an increase in the rates of graft rejection and transplanted organ rejection [12]. Estimates indicate that half of the organ rejections and 15% of the transplant rejections are due to the lack of acceptance and continued use of immunosuppressive medicine [13, 14]. Since even small changes in the immunosuppressive regimen may be associated with poor outcomes and failure in transplantations, timely patient training is of great importance. Given that no adherence to medications is a complex and multifactorial problem, a combination of approaches is highly recommended targeting at different risk factors and building collaboration between patients and medical teams [15, 16].

To this end, most pre- and post-transplant care services usually take place in advanced centers far away from patients causing transportation, social, and economic challenges for the patients and their families [17–19]. The term 'mHealth' refers to the use of smartphones or digital devices for training, managing, monitoring, diagnosing, and treating diseases [20].

During the last decade, advancements in using computers, telemedicine, and mobile health (mHealth) applications, especially in drug evaluation, have affected many areas of healthcare, including transplantation [21]. On the one hand, the adoption of mHealth solutions

is critical in providing knowledge and information to patients and health care providers. On the other hand, applications based on mobile phones are suitable due to their ease of access, acceptability, and support for social distancing, especially during the current COVID-19 epidemic [18, 22]. They also provide innovative solutions for disease management, hope for patients, and help for caregivers and health professionals. These applications cover a wide range of clinical knowledge that can be used by healthcare professionals for specific diseases/disorders in patient-centered approaches [23, 24].

Most of the published telemedicine literature in the field of liver transplant disease is aimed at disease management and improvement of care [25–27]. These scholars noted distance education as a suitable solution for providing patients with advice and supporting them.

The survival rate of patients after receiving a transplant depends on post-transplant care, which requires adequate training of the patient in various fields, such as diet, drug regimen, infectious disease education, daily activities, and physical activities. Education is necessarily needed for liver transplant recipients to promote their health and participate in their own treatment. Educational programs improve health behaviors, effective self-care, symptom management, and control of other diseases caused by liver transplantation [28]. In this regard, information needs analysis and educational contents required by liver transplant recipients are among the main prerequisites for providing proper self-management training by health care providers [11, 29]. As a result, levels of successful transplantation, survival, and quality of life increase. Considering the lack of self-care educational software based on the people's geographical location, race, lifestyle, etc. in Iran, designing and evaluating a mobile phone-based training application seem necessary for liver transplant patients according to the users' needs analysis and opinions of experts. This application can be useful in increasing the patients' levels of awareness and satisfaction.

Methods

Study design

This quantitative study was carried out in two phases. In the first phase, a researcher-made questionnaire was developed and evaluated by a panel of experts who were required to determine the appropriate educational content for the application. In the second phase, the application was designed based on Android and its usability was evaluated by users.

To determine the appropriate educational contents and capabilities required for designing the application program, a list of educational contents and components was prepared followed by reviewing the literature and various

studies in the form of a researcher-made questionnaire. Respondents, including 15 experts (selected based on the purposeful sampling method) were supposed to read the items and evaluate them as necessary (1 score) or unnecessary (0 score). This questionnaire consisted of three sections: participants' demographic information (3 items), educational contents (in 10 general categories and a total of 57 items), and the required capabilities or skills (4 items).

The content validity of the questionnaire was confirmed by six faculty members of medical informatics, gastroenterology, and liver specialty. The questionnaire's reliability was also confirmed by gastroenterologists and liver specialists using the Kuder-Richardson method (KR-20=0.85). After completing the questionnaire, the results were analyzed using descriptive statistics and frequency distribution reports. The educational contents were included in the program in the case that the related items were approved by 65% of the experts.

The application was designed based on the educational content collected from the first phase. This application was developed using Visual Basic programming language, Java Class Library, and in the B4A environment. To evaluate the application's usability, it was installed on the smartphones of Liver transplant patients ($n=30$) referring to Motahari Clinic and Abu Ali Sina Hospital in Shiraz affiliated with Shiraz University of Medical Sciences from May to June 2022. Prior to installing the applications, oral consent was obtained from all participants after providing them with comprehensive explanations of the study goals. Patients were selected based on the number of available cases and their willingness to cooperate in the intervention. Other inclusion criteria were having a smartphone with Android version 5 and higher and having had the transplantation for at least one month. Two weeks after the patients used the application, they were asked to evaluate its usability by the standard Quiz version 5.5 questionnaire. The validity and reliability of the questionnaire have already been confirmed by previous studies. The evaluation section of the questionnaire included: general opinion about using the application (6 items), its screen (4 items), terminology and information (6 items), learnability (6 items), and general capabilities (5 items). Each item was responded based on a 10-point Likert scale from zero (lowest) to nine (highest).

Results

Followed by reviewing the related literature, a questionnaire was developed on educational content required for designing a mobile-based application for patients with liver transplantation. The questionnaire was administered to 15 gastroenterologists and liver specialists who were required to determine the necessary concepts to

be covered by the application. The experts' demographic characteristics (Table 1) showed that most participants were male, were within the age range of 40 to 50 years, and had more than 10 years of work experience.

The participants were asked to read the questionnaire's items carefully and determine the educational content required for developing the application program. Table 2 represents the experts' viewpoints in this vein.

As illustrated in Table 2, the obtained educational components were classified under 10 categories include: transplantation medicine (6 items), nutrition (10 items), exercise (5 items), skin cancer (3 items), cytomegalovirus (4 items), gastrointestinal complications after liver transplant (11 items), diabetes (6 items), oral and dental care (4 items), motherhood (5 items), and sexual desire (3 items). The majority of items were approved by the experts and included in the application. However, liquids (in the nutrition category) and types of exercise (in the activities category) were deemed unnecessary by the experts. The experts' opinions about the application capabilities are listed in Table 3.

An open-ended question was also added to the end of the questionnaire asking the participants to add their possible opinions and suggestions not covered in the questionnaire. After collecting and summarizing the information, 50% of the experts believed that "body mass index (BMI) calculation" and "medication reminders" were among the most significant features of the application design.

Followed by determining the required educational items, a mobile phone-based application was designed and installed on the mobile phones of 30 liver transplant patients who referred to Motahari Clinic and Abu Ali Sina Transplantation Hospital of Shiraz affiliated with Shiraz University of Medical Sciences. To observe ethical considerations, verbal consent was obtained from all patients prior to the intervention.

Table 1 Frequency distribution of experts' demographic characteristics

Variable	n	%
Gender		
Male	12	80
Female	3	20
Age		
< 40	1	6.7
40–50	10	67
> 50	4	26.3
Work Experience		
5–10	1	6.7
11–20	9	60
> 20	5	33.3

Table 2 Frequency distribution of experts' responses regarding the necessity of educational items

Row	Category	Educational items	AGREE	Row	Category	Educational items	AGREE
1	Transplantation medicine	Importance of transplantation medicine	100	29	Gastrointestinal complications after liver transplantation	Digestion of food	67
2		Cyclosporine	100	30		Causes of digestive disorders	93
3		Tacrolimus	100	31		Treatment of gastrointestinal complications	73.4
4		Mycophenolate mofetil	100	32		diarrhea	67
5		Prednisolone and azathioprine	100	33		Symptoms of peptic ulcer	73.4
6		Important points in the using medicine	93	34		Prevention of wounds	93
7	Diet	Fats	100	35	Inflammation of the digestive tract	80	
8		Carbohydrates	100	36	Swallowing stimulants and spicy substances	67	
9		Vitamins	100	37	Esophagitis in transplant recipients	80	
10		Minerals	86.7	38	Gastritis	73.4	
11		Proteins	86.7	39	Pancreatitis	73.4	
12		Fruits and vegetables	80	40	Diabetes	About diabetes	100
13	Milk and diaries	80	41	Post-transplant diabetes		100	
14	Oils	80	42	Reduced risk of diabetes		93	
15	Liquids	60	43	Signs and symptoms of diabetes after transplantation		93	
16	Important recommendations	86.7	44	Complications of Diabetes		100	
17	Activities	Kinds of sports	53.4	45		Living with diabetes	93
18		Starting exercise after transplantation	86.7	46	Oral care	Oral care	86.7
19		Increasing the level of activity	86.7	47		The most common dental problems in transplant recipients	93
20		Appropriate exercises	100	48	Correct brushing	80	
21	Important recommendations	93	49	Performing dental treatments after transplantation	67		
22	Skin Cancer	Skin cancer cases	80	50	Motherhood	Transplantation and pregnancy	93
23		Effective factors	73.4	51		The best time of pregnancy	100
24	Cytomegalovirus	Treatment	73.4	52	Pregnancy complications	93	
25		About cytomegalovirus	86.7	53	Pregnancy risks for the fetus	86.7	
26		Diagnosis of cytomegalovirus	93	54	Lactation	93	
27		Symptoms of cytomegalovirus	93	55	Sexual desires	Sexual desires	93
28	Treatment of cytomegalovirus	86.7	56	Initiation of sexual activity after transplantation		93	
				57		Decreased libido	93

Demographic information of the participants included gender, age, education level, and Duration after the transplant operation (Table 4).

The majorities of participants were Male, had since their transplant for more than a year, and were in the age

range of 40 to 50 years. Considering their level of education, about 66.7% ($n=20$) had a Master's degree or higher. The users' evaluations and

opinions regarding usability of the designed application were collected after two weeks (Table 5).

Table 3 Distribution of the experts' answers regarding the application's capabilities

Capabilities	Necessary	
	n	%
Customizability For the user	12	80
Menus for quick access to content	12	80
The ability to play videos and animations by the user	13	86.7
Introducing digital education resources to the user	14	93.3

Table 4 Frequency distribution of the participants' personal characteristics

Variable	n	%
Age		
< 40	9	30
40–50	17	56.7
> 50	4	13.3
Gender		
Male	22	73.3
Female	8	26.7
Duration after the transplant operation		
Less than a year	10	33.3
1–2 years	13	43.3
More than 2 years	7	23.4
Education level		
High school	2	6.7
Diploma	5	16.6
Associate Degree	3	10
Masters	8	26.7
Masters and above	12	40

Table 5 The end-users' evaluation of the usability of the application

Scale	Mean	SD
Overall Reactions To The Software	8.1	0.64
Screen	7.6	0.93
Terminology And System Information	8.45	0.41
Learning	7.9	0.96
System Capabilities	7.5	0.84
Usability And User Interface	7.9	0.75

Discussion

Given the current need to increase patients' self-care awareness, especially those with chronic diseases, mHealth applications can be employed to reduce costs and enhance the availability of treatment services [30–32]. The present study aimed at designing and evaluating

an educational application for liver transplant patients. To this end, the necessary parameters and educational contents were classified under 10 categories of transplantation medicine, nutrition, daily activities, gastrointestinal complications after liver transplantation, diabetes, skin cancer, megalovirus, oral care, motherhood, and sexual desire were obtained based on the library studies and experts' opinions. The evaluation results indicated that the application was at a good level of usability with an average score of 7.9 ± 0.75 .

In this regard, Wadhvani et al. [33]. Mentioned the medical and social needs of liver transplant recipient children and noted some solutions to improve the care process. The participants, who were parents and caregivers of children, were asked to answer questions about the liver transplant recipient children in an interview. According to the findings, the caregivers were at high levels of motivation and ability regarding the transplantation. The participants also provided the research team with several suggestions to improve care, such as increasing the awareness of parents and children. These researchers concluded that the caregivers' knowledge was not sufficient about post transplantation caring. Consequently, the present study was conducted to increase the awareness of parents and families with Liver transplant recipients.

Joyce et al. [34] carried out a six-month educational course to empower the patients using virtual education. As they reported, the intervention not only obtained positive feedback from the patients but also increased their awareness and participation. The participants maintained the significant role of technology, better support, development of a standard workflow, etc. Our study is in line with this study except that we administered the training program based on a mobile application and tried our best to achieve a high level of usability based on the end-users' needs.

Considering the liver transplant recipients' rapid weight gain and lack of exercise, Hickman et al. [35]. Applied remote health to change their lifestyle (including exercise and diet). Through a 12-week intervention via video telecast, 14 sessions were held by nutrition and exercise specialists for liver transplant recipients. Similar to our study, the post-intervention findings confirmed high levels of reliability and effectiveness for the program, reduced severity scores of metabolic syndromes, and improved diet adherence, quality of life, and mental health. However, all the required educational content and components (including nutrition) were collected through a literature review, corroborated by the experts via a questionnaire, administered among patients (as end users) via an application based on a mobile phone platform, and evaluated in terms of its usability.

Gordon et al. [36]. Evaluated the effectiveness of a web-based application entitled “Notify Me” to increase awareness among kidney recipients. Participants were randomly assigned to the control (regular training) and intervention (regular training along with using the “Notify Me” application) groups. The evaluation was performed in two phases using questionnaires and phone calls. A comparison between scores of the two study groups showed that members of the intervention group obtained a higher knowledge score after the intervention. In another study, Korus et al. [37]. Investigated the acceptance and effectiveness of a web-based self-care application designed for adolescent liver transplant candidates or recipients. The control group had no access to the application while the intervention group members were required to use the application. The evaluation criteria included the application’s feasibility and usage rate, health-related outcomes in self-care assessment, and knowledge level. Data were collected using a semi-structured interview after the intervention and the findings represented higher levels of motivation to practice and learn in the intervention group and patients who had received the transplantation less than a year before the data collection. The participants mentioned obstacles, such as slowness and lack of constant access to the program. As these researchers found, no significant difference was observed between the groups with regard to health-related outcomes. In the same vein, Wedd et al [38]. Investigated the use of portals in liver and kidney transplant recipients ($n=710$) according to the patients’ specific needs at the Transplantation Center of Southeastern, United States. The portal had features, such as drug lists, laboratory results, secure messages, health records, etc. The findings showed the significant effect of educational conditions and race on the users’ application of the portal. Due to the portal’s efficiency, it has been recommended as a suitable tool for health management and patient support. In comparison with the above-mentioned studies, the present study employed a panel of experts to determine the appropriate training content, designed an application based on mobile phones, and evaluated usability of the application.

In a prospective cohort study, Koc et al. [39]. Examined the clinical feasibility, safety, and beneficial effects of a telemedicine-based tele monitoring program (TRMP) for continuous follow-up of adult liver transplant recipients. This study was conducted at the University Hospitals of Leuven, Belgium. The findings did not indicate any increase in the rate of liver transplant rejection and the need for hospitalization caused by tele monitoring. Researchers reported the patients’ high levels of participation and willingness as well as stable clinical condition to run TRMP. This program

could reduce the rate of outpatient visits and improve a safe approach. In other words, remote monitoring has provided the opportunity for accurate follow-up of tacrolimus levels. Unlike this study, we conducted an educational program based on the previous studies and experts’ opinions, tried to obtain all the important educational content, and provided the liver transplant recipients with these achievements in the form of an educational program.

In promoting the caring process for patients with chronic diseases, Hayward et al. [40]. Developed a questionnaire and administered it to cirrhosis patients to determine their level of information about liver disease and the required self-management measures. Half of the patients remembered the written information provided by the clinical expert and 64% reported that obtained the information personally, mostly from the Internet. The educational pamphlet administered in this study had a simple structure, descriptive photos, and plain language to provide patients with their needed information between each follow-up period. Different from the mentioned study, we collected and represented all the post-transplantation information up to the time of recovery when patients returned to their previous lifestyle.

Among the limitations of this study, the following can be mentioned. Due to the epidemic of COVID-19 during the study period, the specialists participating in this research were selected from one of the specialized liver transplant hospitals in Iran. In some cases, experts did not complete the questionnaires comprehensively due to their busy schedules and/or personal lack of motivation. To meet this lack of cooperation by experts, the researcher referred back to the specialist with an incomplete questionnaire, explained the study’s significance and purpose, and asked them to complete the questionnaire. Also, due to the aforementioned problems in accessing the participants, it was not possible to use a questionnaire for nutritional risk, so only parameters such as BMI were used. In this study, the questionnaire’s evaluation section only consisted of the application’s usability. Moreover, only the experts determined and confirmed the educational content needed for the application. To obtain more generalizable results, future researchers are suggested to include experts from the specialized transplantation centers in Iran and obtain the end users’ points of view along with the experts’ ideas to determine the required educational content.

Conclusion

The present study aimed to design an educational mobile-based application for Liver transplant patients based on Android. As a result, the necessary educational content and required educational components were obtained

and classified under 10 main categories using the experts' points of view. The participants' high evaluation scores showed that users were satisfied with the usability of this program. Regardless of the time and place limitations, this application could increase the patients' awareness in the fields related to their disease, participation and involvement in their treatment process, and quality of care. Furthermore, this application was able to reduce complications of liver transplantation while increasing the public's awareness.

Acknowledgements

Authors appreciate nice cooperation of the experts who participated in this research and the staff of Motahari liver clinic and AbuAli Sina Hospital in Shiraz.

Author's contributions

Mostafa Langarizadeh: Conceptualization, Methodology, Validation, Writing - Review & Editing. Fateme Moghbeli: Conceptualization, Resources, Supervision, Methodology, Visualization. Shamim Ahmadi: Software, Methodology, Conceptualization, Validation, Visualization, Resources, Writing - Review & Editing. Mohammad Hossein Langarizadeh: Data curation, Writing - Review & Editing. Mohammadjavad Sayadi: Data curation, Formal analysis, Validation, Writing - Review & Editing. Fatemeh Sarpourian: Data curation, Validation, Writing - Review & Editing. Seyed Ali Fatemi Aghda: Conceptualization, Methodology, Software, Writing - Original Draft, Visualization, Editing, Project administration. All authors have provided final approval of the version submitted.

Funding

The author(s) received no financial support for the research, authorship, or/and publication of this article.

Availability of data and materials

The data used and analysed during the current study are not publicly available due Iran University of Medical Sciences policy but are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

This study was reviewed and approved by the review board and the ethics committee of Iran University of Medical Sciences (IR.IUMS.REC.1397.017). In this study the Informed consent was obtained from all individual participants included in the study. All methods and protocols were carried out in accordance with relevant guidelines and regulations.

Consent for publication

Not Applicable.

Competing interests

The authors declare that they have no competing interests.

Received: 21 November 2022 Accepted: 30 August 2023

Published online: 09 September 2023

References

- Sharma A, Nagalli S. Chronic liver Disease. StatPearls, Treasure Island (FL); 2021.
- Seto W-K, Mandell MS. Chronic liver disease: global perspectives and future challenges to delivering quality health care. *PLoS ONE*. 2021;16(1):e0243607. <https://doi.org/10.1371/journal.pone.0243607>.
- Raeisi Shahraki H, Pourahmad S, Ayatollahi SMT. Identifying the prognosis factors in death after liver transplantation via adaptive LASSO in Iran. *J Environ Public Health*. 2016;2016:7620157.
- Demir B. Effect of sleep hygiene education on sleep quality in patients with liver transplantation. *Holist Nurs Pract*. 2022;36(5):268–74. <https://doi.org/10.1097/HNP.0000000000000542>.
- Okumura K, et al. Improving Liver Transplant Outcomes for Hepatitis C Virus Hepatocellular Carcinoma in the Direct-Acting Antiviral Therapy Era. In: *Transplantation Proceedings*. Elsevier; 2022.
- Becchetti C, Becchetti C, Dirchwolf M, Banz V, Dufour J-F. Medical management of metabolic and cardiovascular complications after liver transplantation. *World J Gastroenterol*. 2020;26(18):2138–54. <https://doi.org/10.3748/wjg.v26.i18.2138>.
- Langarizadeh M, Langarizadeh M, Nadjarzadeh A, Maghsoudi B, Fatemi Aghda SA. The nutritional content required to design an educational application for infertile women. *BMC Womens Health*. 2023;23(1):1–7. <https://doi.org/10.1186/s12905-023-02156-y>.
- Shahmoradi L, Shahmoradi L, Langarizadeh M, Pourmand G, Fard Z, Borhani A. Comparing three data mining methods to predict kidney transplant survival. *Acta Informatica Medica*. 2016;24(5):322. <https://doi.org/10.5455/aim.2016.24.322-327>.
- Zengin M, Yayan EH, Düken ME. The effects of a therapeutic play/play therapy program on the fear and anxiety levels of hospitalized children after liver transplantation. *J PeriAnesthesia Nurs*. 2021;36(1):81–5.
- Cao C, et al. Employment and patient satisfaction after liver transplantation. *J Clin Translational Hepatol*. 2020;8(3):299.
- Akbulut S, Akbulut S, Ozer A, Saritas H, Yilmaz S. Factors affecting anxiety, depression, and self-care ability in patients who have undergone liver transplantation. *World J Gastroenterol*. 2021;27(40):6967–84. <https://doi.org/10.3748/wjg.v27.i40.6967>.
- Uebayashi EY, Uebayashi EY, Okajima H, Yamamoto M, Ogawa E, Okamoto T, Haga H, Hatano E. The new challenge in pediatric liver transplantation: chronic antibody-mediated rejection. *J Clin Med*. 2022;11(16): 4834. <https://doi.org/10.3390/jcm11164834>.
- Di Maira T, Little EC, Berenguer M. Immunosuppression in liver transplant. *Best Pract Res Clin Gastroenterol*. 2020;46:101681.
- Chen J, Chen J, Ferreira J, Martinez M, Lobritto S, Goldner D, Vittorio J. Role of budesonide for the treatment of rejection in pediatric liver transplantation. *J Pediatr Gastroenterol Nutr*. 2020;71(3):388–92. <https://doi.org/10.1097/MPG.0000000000002784>.
- Ko D, Muehrer RJ, Bratzke LC. Self-management in liver transplant recipients: a narrative review. *Prog Transpl*. 2018;28(2):100–15. <https://doi.org/10.1177/1526924818765814>.
- Bilhartz JL, Bilhartz JL, Lopez MJ, Magee JC, Shieck VL, Eder SJ, Fredericks EM. Assessing allocation of responsibility for health management in pediatric liver transplant recipients. *Pediatr Transplant*. 2015;19(5):538–46. <https://doi.org/10.1111/ptr.12466>.
- Langarizadeh M, Langarizadeh M, Rahmati P, Yousefpour Azari S, Sarpourian F, Sayadi MJ, Langarizadeh MH, Fatemi Aghda SA. Identifying and validating the educational needs to develop a celiac self-care system. *BMC Prim Care*. 2023;24(1):1–8. <https://doi.org/10.1186/s12875-023-02076-8>.
- Melilli E, Melilli E, Cestone G, Revuelta I, Meneghini M, Lladó L, Montero N, Manonelles A, Diaz M, Coloma A, Torregrosa V, Baliellas C, Cruzado JM, Diekmann F, Grinyó J, Bestard O. Adoption of a novel smart mobile-health application technology to track chronic immunosuppression adherence in solid organ transplantation: results of a prospective, observational, multicentre, pilot study. *Clin Transplant*. 2021;35(5): e14278. <https://doi.org/10.1111/ctr.14278>.
- Fallah-Aliabadi S, et al. Hospital disaster resilience: development of an assessment tool using expert panel and fuzzy analytical network process. *Int J Disaster Resil Built Environ*. 2022;13(5):554–67.
- Langarizadeh M, Fatemi Aghda SA, Nadjarzadeh A. Design and evaluation of a mobile-based nutrition education application for infertile women in Iran. *BMC Med Inf Decis Mak*. 2022;22(1):58.
- Langarizadeh M, Fatemi Aghda SA, Nadjarzadeh A. Design and evaluation of a mobile-based nutrition education application for infertile women in Iran. *BMC Med Inf Decis Mak*. 2022;22(1):1–9.
- Marzaleh MA et al. Application of telerehabilitation for older adults during the COVID-19 pandemic: A systematic review. *Disaster Med Pub Health Prep*. 2022;17:e402p.
- Khundaqji H, Khundaqji H, Hing W, Furness J, Climstein M. Wearable technology to inform the prediction and diagnosis of cardiorespiratory events: a scoping review. *PeerJ*. 2021;9: e12598. <https://doi.org/10.7717/peerj.12598>.

24. Samad-Soltani T, Ghanei M, Langarizadeh M. Development of a fuzzy decision support system to determine the severity of obstructive pulmonary in chemical injured victims. *Acta Informatica Medica*. 2015;23(3):138.
25. Le LB, Le LB, Rahal HK, Viramontes MR, Meneses KG, Dong TS, Saab S. Patient satisfaction and healthcare utilization using telemedicine in liver transplant recipients. *Dig Dis Sci*. 2019;64(5):1150–7. <https://doi.org/10.1007/s10620-018-5397-5>.
26. Kröncke S, Kröncke S, Lund LK, Buchholz A, Lang M, Briem-Richter A, Grabhorn EF, Sterneck M. Psychosocial situation, adherence, and utilization of video consultation in young adult long-term pediatric liver transplant recipients during COVID-19 pandemic. *Pediatr Transplant*. 2021;25(8): e14121. <https://doi.org/10.1111/ptr.14121>.
27. Santonicola A, Santonicola A, Zingone F, Camera S, Siniscalchi M, Ciacci C. Telemedicine in the COVID-19 era for liver transplant recipients: an Italian lockdown area experience. *Clin Res Hepatol Gastroenterol*. 2021;45(3):101508. <https://doi.org/10.1016/j.clinre.2020.07.013>.
28. Yang LS, Yang LS, Shan LL, Saxena A, Morris DL. Liver transplantation: a systematic review of long-term quality of life. *Liver Int*. 2014;34(9):1298–313. <https://doi.org/10.1111/liv.12553>.
29. Sá AS et al. Information needs of family caregivers regarding liver transplant candidates. *Rev Gaucha Enferm*. 2016;37(1):e54650.
30. Debon R, et al. Mobile health applications for chronic diseases: a systematic review of features for lifestyle improvement. Volume 13. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*; 2019. pp. 2507–12. 4.
31. Bienfait F, et al. Applying m-Health to palliative care: a systematic review on the use of m-Health in monitoring patients with chronic diseases and its transposition in palliative care. *Am J Hospice Palliat Medicine*. 2020;37(7):549–64.
32. Luo H, et al. An m-Health intervention for rheumatoid arthritis in China (Rheumatism Center app): study protocol for a prospective Randomized Controlled Trial. *Nurs Open*. 2022;9(6):2915–24.
33. Wadhvani S et al. Caregiver perspectives on the everyday medical and social needs of long-term pediatric liver transplant patients. *Liver Transpl*. 2022;28(11):1735–46.
34. Joyce M, et al. Expansion of patient education programming regarding live donor liver transplantation via virtual group encounters during the COVID-19 pandemic. In: *Transplantation Proceedings*. Elsevier; 2021.
35. Hickman IJ et al. Telehealth-delivered, cardioprotective diet and exercise program for liver transplant recipients: a randomized feasibility study. *Transplantation Direct*. 2021;7(3):e667.
36. Gordon EJ, et al. Effect of a mobile web app on kidney transplant candidates' knowledge about increased risk donor kidneys: a randomized controlled trial. *Transplantation*. 2017;101(6):1167.
37. Korus M, Korus M, Cruchley E, Calic M, Gold A, Anthony SJ, Parekh RS, Stinson JN. Assessing the acceptability and efficacy of teens taking charge: Transplant—A pilot randomized control trial. *Pediatr Transplant*. 2020;24(1): e13612. <https://doi.org/10.1111/ptr.13612>.
38. Wedd J, et al. Racial, ethnic, and socioeconomic disparities in web-based patient portal usage among kidney and liver transplant recipients: cross-sectional study. *J Med Internet Res*. 2019;21(4):e11864.
39. Koc ÖM, Koc ÖM, Pierco M, Remans K, Van den Hende T, Verbeek J, Van Malenstein H, Van der Merwe S, Robaey G, Monbaliu D, Pirenne J, Van den Bosch B, Dobbels F, Nevens F. Telemedicine based remote monitoring after liver transplantation: feasible in a select group and a more stringent control of immunosuppression. *Clin Transplant*. 2022;36(1): e14494. <https://doi.org/10.1111/ctr.14494>.
40. Hayward KL, Hayward KL, Horsfall LU, Ruffin BJ, Cottrell WN, Chachay VS, Irvine KM, Martin JH, Powell EE, Valery PC. Optimising care of patients with chronic disease: patient-oriented education may improve disease knowledge and self-management. *Intern Med J*. 2017;47(8):952–5. <https://doi.org/10.1111/imj.13505>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

